

Successful Management With Interferon Alpha-2a After Prednisone Therapy Failure in an Infant With a Giant Cavernous Hemangioma

Manuel A. Castello, MD,^{1*} G. Ragni, MD,¹ A. Antimi, MD,¹ A. Todini, MD,²
G. Patti, MD,³ R. Lubrano, MD,¹ A. Clerico, MD,¹ and A. Calisti, MD³

A giant cavernous hemangioma of the left arm with severe thrombocytopenia and consumptive coagulopathy was observed in a neonate. Initial treatment with prednisone, platelet transfusions, and clotting replacement failed to control the bleedings. The child was then treated with daily

subcutaneous infusions of interferon alpha-2a. Coagulopathy rapidly improved and transfusions were drastically reduced. The hemangioma regressed progressively and disappeared after 4 months of treatment. **Med. Pediatr. Oncol.** 28: 213–215 © 1997 Wiley-Liss, Inc.

Key words: giant cavernous hemangioma; Kasabach-Merritt syndrome; medical therapy

INTRODUCTION

Hemangiomas are common in infancy and usually follow a benign course. The spontaneous regression rate is 80–85% [1]. Rarely, they can be giant, produce skeletal overgrowth or high cardiac output failure, or be associated with severe thrombocytopenia, microangiopathic hemolytic anemia, and bleeding. This latter association was described for the first time in a 2-year-old child by Kasabach-Merritt [2] in 1940.

CASE REPORT

A 29-year-old woman (gravidia 1, para 1) gave birth at term to a baby girl weighing 3.050 kg. The Apgar score was 7 after the first minute and 9 after the fifth. Postpartum examination revealed a giant hemangioma extending from the left anterolateral chest wall to the middle third of the forearm (Fig. 1). The limb was maintained in extension, and the overlying skin was bluish and intact. She exhibited hematomas and widespread petechiae over her body. Neither hematuria, nor internal hemorrhages, resulted. The ECG showed right ventricular overload, and the echocardiogram, a mild ventricle enlargement.

Laboratory data revealed a severe consumptive coagulopathy (hemoglobin 7.2 g/dl, platelets 10,000/mm³, prothrombin time >60 sec (n.v. 13–20 sec), partial thromboplastin time 35 sec (n.v. <55 sec), fibrinogen undetectable (n.v. 125–300 mg/dl), unconjugated bilirubin 7.2 mg/dl (n.v. <12 mg/dl), Coombs' tests were negative). Initial management consisted of platelet and red blood cell transfusions, clotting factors replacement, and corticosteroids (prednisone, 7.5 mg daily). During the following 2 weeks, 12 platelet and eight red blood cell transfusions were administered together with clotting factors replacements,

and prednisone 7.5 mg daily, but no significant result was obtained. Her general conditions worsened: the consumptive coagulopathy, thrombocytopenia (platelets 11,000/mm³) and subsequent anemia (Hb 7.1 g/dl) did not subside. The giant hemangioma and consequent edema did not regress, and a large hematoma (maximum diameter 12 cm) was observed over the left shoulder and latero-cervical region.

Interferon alpha-2a therapy was then initiated (3,000,000 U/m²/day subcutaneously). After the first 10 days of treatment, her general conditions rapidly improved (prothrombin time, 18 sec; partial thromboplastin time, 31 sec; fibrinogen, 156 mg/dl; Hb, 8.2 g/dl; platelets, 12,000/mm³). The number of transfusions decreased from 28, administered during the first month of life, to eight during the second.

The hemangioma regressed in size, disappearing from the forearm and chest wall after the first month of treatment (Fig. 2). Transfusions and clotting factors replacement were discontinued during the second month of life when laboratory data revealed progressively increasing values that returned to normal (prothrombin time, 13.1 sec; partial thromboplastin time, 28 sec; fibrinogen, 250 mg/dl; Hb, 13.5 g/dl; platelets, 253,000/mm³) during the following month (Fig. 3). After the fourth month of life, interferon alpha-2a dosage was decreased to 3,000,000 U/m² every other day. The hemangioma continued to

¹Department of Pediatrics, University "La Sapienza," ²Departments of Angiology and ³Pediatric Surgery, Ospedale San Camillo, Rome, Italy.

*Correspondence to: Manuel A. Castello, M.D., Università degli Studi di Roma "La Sapienza," Istituto di Clinica Pediatrica, S.S. Oncologia Pediatrica, V. le Regina Elena 324, 00161 Rome, Italy.

Received 20 February 1996; accepted 6 May 1996.



Fig. 1. Giant hemangioma extending from the left anterolateral chest wall to the middle third of the forearm (middle arm and forearm circumference: 23/14 cm, respectively), at birth.



Fig. 2. The hemangioma regressed in size, disappearing from the forearm and chest wall after the first month of treatment.

regress until complete disappearance after 4 months of treatment. There were no side effects. During this period, the child's growth was adequate. In accordance with the literature [16], interferon alpha-2a was discontinued after 6 months of treatment. At present, the child is well and no relapse has occurred 26 months after complete remission. The left arm is normal (Fig. 4), there are no differences in size, color, or function between the two arms; no skeletal overgrowth occurred. The regression pattern of the hemangioma was followed-up with echo-color Doppler investigations performed every month.

ECHO-COLOR DOPPLER REPORTS

The echo-color Doppler examination is important in the detection of vascular dysplasia, and in distinguishing between venous and artero-venous (A-V) types. Furthermore, with this technique the evolution of the dysplasia can be safely followed-up. In the case reported here, the examination performed on the third day of life showed




Fig. 3. Laboratory data and transfusions vs. treatment during the first 2 months of life. PT (prothrombine time in seconds), T (number of red blood cell and platelet transfusions), prednisone: 7.5 mg/day; interferon alpha 2a: 3,000,000 U/sqm/day.



Fig. 4. At present, the left arm is normal, there are no differences in size, color, or function between the two arms; no skeletal overgrowth occurred.

a giant hemangioma with high and low load A-V shunts located in the anterolateral chest wall, arm, and forearm. Small caliber veins with high velocity flow were also seen. The peripheral venous distribution of the arm was normal. On the fifth day, the situation worsened with the development of edema in the upper limb that compressed the veins. At this point, treatment with interferon was started. Controls performed every month showed a progressive improvement in the ultrasound findings: a progressive reduction of the A-V shunts that gradually disappeared from the chest wall and forearm, an increase in the caliber of veins, and a decrease in venous flow. Finally, the last control revealed a similar arterovenous flow (velocytogram) in the two limbs. Microshunts that had been observed in the proximal region of the cephalic vein, and that had caused an increase in load and venous flow in the cephalic and innominate vein, had also disappeared.

DISCUSSIONS AND CONCLUSIONS

Giant cavernous hemangiomas may cause severe hemorrhages, microangiopathic hemolytic anemia, or signifi-

cant cardiac failure due to arterovenous fistula [2-4]. Various types of therapy for the control and/or cure of these complications have been attempted with different results [5]. Current management of hemangiomas includes drugs (e.g., corticosteroids, aminocaproic acid, and heparin to treat the consumptive coagulopathy, and cyclophosphamide), surgical excision, radiotherapy, and argon laser therapy [6-13]. Surgery is the treatment of choice, but the size and/or anatomic location of the lesion may preclude its removal. Pharmacologic therapy is indicated for inoperable hemangiomas, and for life-threatening complications such as congestive heart failure or bleeding diathesis. Corticosteroids are first-line treatment, and the response rate varies from 30 to 60%.

In patients unresponsive to corticosteroids, treatment is not well defined. A prospective study from the Children's Hospital of Boston evaluated interferon alpha-2a in patients with life-threatening hemangiomas unresponsive to corticosteroids [14-15]. Interferon alpha-2a was used because of its reported inhibition of proliferation of vascular cells [16,17] and minimal side effects [16-20] (e.g., neutropenia, mild fever, and abnormal liver function tests). The response rate was 80% with a mean response time of 7 to 8 months [16]. The pharmacologic effects of interferon alpha-2a remain unclear. It has been shown that endothelial cell proliferation and angiogenesis are inhibited by interferon alpha-2a. The resolution of consumptive coagulation seems related to the decrease in thrombocyte aggregation that may be explained by a reduction in platelet adherence and trapping resulting from the increasing endothelial synthesis of prostacyclin induced by interferon alpha [18-20]. In the case reported here, the reduction in platelet and blood transfusions, the progressive involution in size, and the gradual normalization of the laboratory test values, seem strongly related to interferon administration.

ACKNOWLEDGMENT

This work was supported by CNR ACRO 94.01110.39.

REFERENCES

1. Edgerton MT, Morgan R: Hemangiomas: congenital hamartomas. In Welch KJ, Randolph JG, Ravitch MM, O'Neill JA Jr, Rowe MI (eds): "Pediatric Surgery." 4th ed. vol 2. Chicago: Year Book Medical Publishers, Inc. 1986, pp. 1511-1518.
2. Kasabach HH, Merritt KK: Capillary haemangiomas with extensive purpura. *Am J Dis Child* 59:1063-1070, 1940.
3. Ciardini A, Binio G, Gabbiani A: Contributo clinico alla sindrome di Kasabach-Merritt. *Minerva Pediatr* 29:1017-1020, 1977.
4. Currie BG, Schell D, Bowring AC: Giant hemangioma of the arm associated with cardiac failure and the Kasabach-Merritt syndrome in a neonate. *J Pediatr Surg* 6:734-737, 1991.
5. Larsen E, Zinkham W, Eggleston J, Basil JZ: Kasabach-Merritt syndrome: therapeutic considerations. *Pediatrics* 6:971-979, 1987.
6. Corby DG, Oquendo-Cabrera A, Louro JM: Thrombocytopenia in a patient with a relatively small facial hemangioma. Successful management with irradiation after failure of prednisone therapy. *Clin Pediatr* 12:728-731, 1969.
7. Macey RF, Camisa C: Kasabach-Merritt syndrome. *Pediatr Dermatol* 2:133-136, 1991.
8. Nakada K, Kawada T, Fujoka T, Kitagawa H, Enami T, Kuwabara M, Kawaguchi F, Wakisawa M, Yamate M: Hemangioma of the upper arm associated with massive hemorrhage in a neonate. *JPN Surg* 23:273-276, 1993.
9. Vecchi V, Faldella G, Putti C: Sindrome di Kasabach-Merritt associata a coagulazione intravascolare disseminata. *Minerva Pediatr* 29:2335-2339, 1977.
10. Miller JC, Orton CI: Long term follow-up of a case of Kasabach-Merritt syndrome successfully treated with radiotherapy and corticosteroids. *Brit J Plas Surg* 45:559-561, 1992.
11. El-Dessouky M, Azmy AF, Raine PAM, Young DJ: Kasabach-Merritt syndrome. *J Pediatr Surg* 2:109-111, 1988.
12. Tan OT: Laser for vascular lesions in pediatric dermatology. *Pediatr Dermatol* 9:358-360, 1992.
13. Neidhart J, Roach R: Successful treatment of skeletal hemangioma and Kasabach-Merritt syndrome with aminocaproic acid. Is fibrinolysis defensive? *Am J Med* 73:434-437, 1982.
14. Mulliken JB: A biological approach to cutaneous vascular anomalies. *Pediatr Dermatol* 9:356-357, 1982.
15. Brouty-Boye D, Zetter BR: Inhibition of cell motility by interferon. *Science* 208:516-518, 1980.
16. Hatley RM, Sabio M, Howell CG: Successful management of an infant with a giant hemangioma of the retroperitoneum and Kasabach-Merritt syndrome with alpha interferon. *J Pediatr Surg* 28:1356-1359, 1993.
17. Klein C, Hauser M, Hadorn HB: Interferon alpha-2a therapy of consumptive coagulopathy in Kasabach-Merritt syndrome. *N Engl J Med* 327:919, 1992.
18. Orchard PJ, Smith C, Woods WG, Day DL, Dehner LP, Shapiro R: Treatment of haemangioendotheliomas with alpha interferon. *Lancet* 2:565-567, 1989.
19. Ezekowitz A, Mulliken J, Folkman J: Interferon alpha therapy of haemangiomas in newborn and infants. *Br J Hematol* 79:67-68, 1991.
20. White C, Sondheimer HM, Crouch E, Wilson H, Leland LF: Treatment of pulmonary hemangiomatosis with recombinant interferon alpha-2a. *N Engl J Med* 18:1197-1200, 1989.